

Comment on “Radiative corrections and parity nonconservation in heavy atoms”

M. Yu. Kuchiev* and V. V. Flambaum†

School of Physics, University of New South Wales, Sydney 2052, Australia

(Dated: February 1, 2008)

PACS numbers: 11.30.Er, 31.30.Jv, 32.80.Ys

A recent paper of Milstein *et al* [1], which is commented below, presents an opportunity to summarize briefly interesting results that have been obtained for QED corrections to the parity nonconservation amplitude (PNC) in atoms.

It has become clear quite recently, that the QED radiative corrections to the PNC amplitude in atoms are sufficiently large, of the order of $\sim 1\%$, and therefore should be included in an analysis of experimental data. High accuracy of this data (0.3% for the $6s - 7s$ PNC amplitude in ^{133}Cs [2]) is used as a test of the standard model, prompting accurate treatment of the radiative corrections. The most difficult for calculations (and even estimations) part of the QED corrections presented the self-energy and vertex corrections to the PNC amplitude. They also prove to give the largest contribution, as has become clear only during last several months, while beforehand they were assumed to be negligible small. It should be noted that without the self-energy corrections the experimental data of Ref. [2] deviate by more than 2σ from predictions of the standard model, for a brief guide to the modern literature devoted to the subject see [3]. Remarkably, when these corrections are calculated accurately, their contribution brings the experimental results of [2] in full agreement with the standard model, as was first demonstrated in [3] and confirmed by recent [4] and [1]. The above mentioned facts have made last developments in the outlined research area quite interesting and exciting.

Fig. 1 presents a summary of results for the QED self-energy (plus vertex) radiative corrections to the PNC amplitude in atoms. It shows these corrections in relative units, as a percent of the PNC amplitude, versus the charge of the atomic nucleus considered. The most interesting for recent experimental applications atoms are Cs, $Z = 55$, and Tl, Pb and Bi with Z close to 80. Fig. 1 shows a close agreement of results of our Ref. [3] with the more recent results of [1, 4]. The important conclusion that follows from this figure is that the self-energy correction to the PNC amplitude is negative and large. For the Cs atom there exists perfect agreement between all the results $-0.73(20)\%$ [3], $-0.9(2)\%$ [4], and -0.85% [1]. A slight discrepancy observed for more heavy atoms Tl, Pb, and Bi, -1.6% [3, 4] and -1.4% [1], is, probably, within the errors of the methods used, as is specified below. As was demonstrated in [3] and supported by [1, 4], the found correction brings the experimental data




FIG. 1: QED self-energy radiative corrections (%) for the PNC amplitude in atoms versus the atomic nuclear charge Z . The thick short-dashed line, thick solid line and thin solid line - results of papers [3], [4], and [1] respectively. The thin dashed line - the lowest term of αZ expansion $-1.97\alpha^2 Z$ [1, 4].

of Wood *et al* [2] within the limits of the standard model.

A reliability of this conclusion is supported by two different groups which have converged to one and the same result. This positive development is further magnified by the fact that previously there has existed a sharp contradiction between [3] and the results of Milstein and Sushkov who insisted that the self-energy correction is small and positive [5], casting doubts on possible relation between experimental data of [2] and the standard model.

Note that the results presented in Fig. 1 were obtained in [1, 3, 4] using different methods and approximations. The work [3] takes into account all orders in the αZ -expansion deriving an identity which relates the radiative corrections to the PNC matrix elements between the $s_{1/2}$ and $p_{1/2}$ electron wave functions with the radiative corrections to the finite nuclear size (FNS) energy shifts of the $s_{1/2}$ and $p_{1/2}$ levels. The works [1, 4] calculated the self-energy corrections to PNC using the αZ -expansion. The results of these latter works can be used to test validity of the identity found in Ref. [3]. Numerical results of [1, 4] shown in Fig. 1 indicate that this identity holds, bringing the final conclusions of these works in very close agreement with [3]. (Note that a possible $\leq 20\%$ violation of the identity, that was proclaimed in [1], is smaller than the error that follows from the parameter of the per-

turbative expansion used in [1, 4], which is $\alpha Z = 0.4$ for Cs and $\alpha Z = 0.6$ for Tl, Pb, Bi. The mentioned deviation appeared probably because the $p_{1/2}$ FNS was extrapolated in [1] to small- Z values from the high- Z FNS of the $2p_{1/2}$ state calculated numerically in different works - see Refs. in [1, 3]. This extrapolation should incorporate an error governed by the parameter mentioned above [6].) The work [1] calculates also corrections $\sim \alpha^3 Z^2 \ln mR$ where R is the nuclear radius verifying (in this order) that the identity of [3] holds exactly, as it should.

Thus, the results obtained by completely different methods agree. We believe this signifies a point after which the discrepancy between the atomic experimental data of [2] and the standard model disappears.

This work was supported by the Australian Research Council.

* email: kuchiev@newt.phys.unsw.edu.au

† email: flambaum@newt.phys.unsw.edu.au

- [1] A.I.Milstein, O.P.Sushkov, and I.S.Terekhov, hep-ph/0208227 (2001).
- [2] C.S.Wood, S.C.Bennett, D.Cho, B.P.Masterson, J.L.Roberts, C.E.Tanner, and C.E.Wieman, Science 275, 1759 (1997).
- [3] M.Yu.Kuchiev and V.V.Flambaum, hep-ph/0206124 (2002).

- [4] M.Yu.Kuchiev, hep-ph/0208196 (2002).
- [5] A.I.Milstein and O.P.Sushkov, Phys. Rev. **66** 022108-1 (2002); hep-ph/0109257 (2001).
- [6] Milstein *et al* [1] also expressed their concerns about our derivation of the identity [3] that used a particular gauge transformation with several gauge parameters. We believe the derivation is correct, but since it proved to be uncomfortable for the authors of [1] we can modify the derivation using the gauge transformation with only *one* parameter. The only condition that is vital for this derivation is the one that states that the vertex contribution to the PNC amplitude equals half the sum of the vertex contributions to the FNS energy shifts of the $s_{1/2}$ and $p_{1/2}$ levels (all corrections in relative units). If this condition holds, then the vertex correction does not contribute to our identity, while the self-energy contributions are taken care of as is specified in [3]. To satisfy the necessary condition we make the gauge transformation with only one gauge parameter. Alternatively, one can resort to a simplified, approximate derivation that does not rely on gauge transformations at all. One neglects the electron mass (which is not important at small distances) in the vertex diagram and commutes the Dirac matrix γ_5 from the weak interaction vertex to the side. Then one uses a relation for the electron wave functions at small distances $\gamma_5 \psi_s \propto \psi_p$. As a result the vertex diagram for the PNC amplitude is transformed into the vertex diagram for the FNS energy shift of either $p_{1/2}$ or $s_{1/2}$ level. An error introduced by this simplified derivation does not exceed the difference in relative corrections to the FNS energy shifts for the considered $p_{1/2}$ and $s_{1/2}$ levels.